

County: Sample Kansas State Total indicators: 124 Tuesday 14th of July 2015 09:30:17 AM

| | County Di | stribution | Value | | Target | | | |
|---|------------|------------|----------|------------|--------------|------------|--------------|------------------|
| ndicator | State | US | State | US | HP2020 Local | Trend | Score | Precision |
| /orkers who Walk to Work | 3 | 3 | 3 | 3 | 3 | 2 | 2.78 | High |
| epression: Medicare Population | 3 | 3 | 2 | 3 | 1.5 | 3 | 2.67 | High |
| /orkers who Drive Alone to Work | 3 | 3 | 2 | 3 | 1.5 | 3 | 2.67 | High |
| ouseholds with Cash Public Assistance Income | 3 | 3 | 3 | 3 | 1.5 | 2 | 2.61 | High |
| hronic Kidney Disease: Medicare Population | 3 | 2 | 3 | 2 | 1.5 | 3 | 2.50 | High |
| tudents Eligible for the Free Lunch Program | 3 | 2 | 3 | 1.5 | 1.5 | 3 | 2.42 | Medium |
| overty Status by School Enrollment | 2 | 2 | 3 | 2 | 1.5 | 3 | 2.33 | High |
| ge-Adjusted Years of Potential Life Lost - Alzheimers | 3 | 1.5 | 2 | 1.5 | 1.5 | 3 | 2.17 | Medium |
| ge-Adjusted Years of Potential Life Lost - Diabetes | 2 | 1.5 | 3 | 1.5 | 1.5 | 3 | 2.17 | Medium |
| hildren Living Below Poverty Level | 2 | 1 | 3 | 2 | 1.5 | 3 | 2.17 | High |
| omeownership | 3 | 2 | 2 | 1 2 | 1.5 | 3 | 2.17 | High |
| oung Children Living Below Poverty Level | 3 | 3 | 3 1.5 | 1.5 | 1.5 1.5 | 3 2 | 2.17 2.11 | High |
| rocery Store Density exually Transmitted Disease Rate | 3 | 1.5 | 3 | 1.5 | 1.5 | 2 | 2.11 | Medium Medium |
| tudent-to-Teacher Ratio | 3 | 3 | 2 | 1.5 | 1.5 | 1.5 | 2.11 | Medium |
| eople 65+ Living Alone | 2 | 3 | 2 | 3 | 1.5 | 1.5 | 2.06 | High |
| ast Food Restaurant Density | 3 | 3 | 1.5 | 1.5 | 1.5 | 1.5 | 2.00 | Low |
| eople Living Below Poverty Level | 2 | 1 | 3 | 1 | 1.5 | 3 | 2.00 | High |
| NAP Certified Stores | 3 | 3 | 1.5 | 1.5 | 1.5 | 1.5 | 2.00 | Medium |
| ge-adjusted Chronic Lower Respiratory Disease Mortality Rate per 100 | 2 | 1.5 | 3 | 1.5 | 1.5 | 2 | 1.94 | Medium |
| ge-adjusted Unintentional Injuries Mortality Rate per 100,000 populat | 1 | 1.5 | 3 | 1.5 | 3 | 2 | 1.94 | Medium |
| ge-Adjusted Years of Potential Life Lost - Chronic Lower Respiratory Dis | 2 | 1.5 | 3 | 1.5 | 1.5 | 2 | 1.94 | Medium |
| ercent of Births with Low Birth Weight | 3 | 1.5 | 3 | 1.5 | 2 | 1 | 1.94 | Medium |
| armers Market Density | 2 | 2 | 1.5 | 3 | 1.5 | 1.5 | 1.92 | Medium |
| ood Environment Index | 3 | 2 | 2 | 1.5 | 1.5 | 1.5 | 1.92 | Medium |
| ercent of Adults Who Currently Smoke Cigarettes | 1.5 | 1.5 | 3 | 1.5 | 3 | 1.5 | 1.92 | Low |
| evere Housing Problems | 3 | 2 | 2 | 1.5 | 1.5 | 1.5 | 1.92 | Medium |
| ge-Adjusted Years of Potential Life Lost - Homicide | 3 | 1.5 | 3 | 1.5 | 1.5 | 1 | 1.89 | Medium |
| Izheimer's Disease or Dementia: Medicare Population | 2 | 3 | 2 | 2 | 1.5 | 1 | 1.89 | High |
| nfant Mortality Rate | 2 | 1.5 | 3 | 1.5 | 3 | 1 | 1.89 | Medium |
| ife Expectancy for Females | 3 | 2 | 2 | 2 | 1.5 | 1 | 1.89 | High |
| ercent of Births Occurring to Unmarried Women | 3 | 1.5 | 3 | 1.5 | 1.5 | 1 | 1.89 | Medium |
| ercent of WIC Mothers Breastfeeding Exclusively | 3 | 1.5 | 3 | 1.5 | 1.5 | 1 | 1.89 | Medium |
| ate of Violent Crime per 1,000 Population | 3 | 1.5 | 3 | 1.5 | 1.5 | 1 | 1.89 | Medium |
| Ininsured Adult Population Rate | 2 | 1.5 1.5 | 3 | 1.5 1.5 | 3 | 1 | 1.89 | Medium |
| ge-Adjusted Years of Potential Life Lost - Heart Disease | 2 | 3 | 3 1.5 | 1.5 | 1.5 1.5 | 1.5 1.5 | 1.83 1.83 | Medium Low |
| hildren with Low Access to a Grocery Store eath Rate due to Drug Poisoning | 1.5 | 2 | 3 | 1.5 | 1.5 | 1.5 | 1.83 | Low |
| viabetes: Medicare Population | 2 | 1 | 2 | 1.5 | 1.5 | 3 | 1.83 | High |
| ood Insecurity Rate | 3 | 2 | 2 | 1 | 1.5 | 1.5 | 1.83 | Medium |
| ercent of Infants Fully Immunized at 24 Months | 2 | 1.5 | 1 | 1.5 | 1.5 | 3 | 1.83 | Medium |
| ercentage of Premature Births | 2 | 1.5 | 2 | 1.5 | 0 | 3 | 1.83 | Medium |
| ge-Adjusted Years of Potential Life Lost - Suicide | 2 | 1.5 | 2 | 1.5 | 1.5 | 2 | 1.78 | Medium |
| amilies Living Below Poverty Level | 2 | 1 | 3 | 1 | 1.5 | 2 | 1.78 | High |
| lomeowner Vacancy Rate | 2 | 2 | 2 | 1 | 1.5 | 2 | 1.78 | High |
| Icohol-Impaired Driving Deaths | 2 | 2 | 2 | 1.5 | 1.5 | 1.5 | 1.75 | Medium |
| ercent of Adults Tested and Diagnosed with High Cholesterol | 1.5 | 1.5 | 2 | 1.5 | 3 | 1.5 | 1.75 | Low |
| ercent of Adults with Diagnosed Diabetes | 1.5 | 1.5 | 3 | 1.5 | 1.5 | 1.5 | 1.75 | Low |
| ercent of Adults with Diagnosed Hypertension | 1.5 | 1.5 | 2 | 1.5 | 3 | 1.5 | 1.75 | Low |
| ge-adjusted Homicide Mortality Rate per 100,000 population | 3 | 1.5 | 3 | 1.5 | 0 | 1 | 1.72 | Medium |
| ge-adjusted Nephritis, Nephrotic Syndrome, Nephrosis Mortality Rate | 2 | 1.5 | 3 | 1.5 | 1.5 | 1 | 1.72 | Medium |
| ge-Adjusted Years of Potential Life Lost - Cancer | 2 | 1.5 | 3 | 1.5 | 1.5 | 1 | 1.72 | Medium |
| louseholds without a Vehicle | 3 | 2 | 3 | 0 | 1.5 | 1 | 1.72 | High |
| ge-adjusted Suicide Mortality Rate per 100,000 population | 1.5 | 1.5 | 1 | 1.5 | 3 | 2 | 1.69 | Medium |
| ge-Adjusted Years of Potential Life Lost - Nephritis, Nephrotic Syndrom | 3 | 1.5 | 3 | 1.5 | 1.5 | 0 | 1.67 | Medium |
| hild Food Insecurity Rate | 2 | 1 | 2 | 2 | 1.5 | 1.5 | 1.67 | Medium |
| ow-Income and Low Access to a Grocery Store | 2 | 2 | 1.5 | 1.5 | 1.5 | 1.5 | 1.67 | Low |
| ublic Water Supply - Percent of Population Served Unaffected by SDW/ | 3 | 1.5 | 1 | 1.5 | 1.5 | 1.5 | 1.67 | Medium |
| nemployed Workers in Civilian Labor Force | 3 | 1 | 3 | 0 | 1.5 | 1.5 | 1.67 | High |
| ercent of Adults who are Obese | 1.5 | 1.5 | 2 | 1.5 | 2 | 1.5 | 1.64 | Low |
| ge-adjusted Cancer Mortality Rate per 100,000 population | 2 | 1.5 | 2 | 1.5 | 2 | 1 | 1.61 | Medium |
| eople Living 200% Above Poverty Level | 1 | 1 | 2 | 2 | 1.5 | 2 | 1.61 | High |
| /orkers Commuting by Public Transportation | 1.5 | 1 | 1 2 | 3 1 E | 3 | | 1.61 | High |
| ercent of Adults Ages 65 Years and Older Who Were Immunized Again: ercent of Adults Who Reported Consuming Fruit Less than 1 Time Per I | 1.5 1.5 | 1.5 1.5 | 2 | 1.5 1.5 | 1.5 1.5 | 1.5 1.5 | 1.58 1.58 | Low Low |
| ercent of Adults Who Were Ever Diagnosed with a Depressive Disorder | 1.5 | 1.5 | 2 | 1.5 | 1.5 | 1.5 | 1.58 | Low |
| ercent of Adults with Fair or Poor Self-Perceived Health Status | 1.5 | 1.5 | 2 | 1.5 | 1.5 | 1.5 | 1.58 | Low |
| ligh School Graduation | 3 | 1.5 | 2 | 1.5 | 2 | 0 | 1.56 | Medium |
| iquor Store Density | 1 | 3 | 0 | 3 | 1.5 | 1 | 1.56 | High |



County: Sample Kansas State Total indicators: 124 Tuesday 14th of July 2015 09:30:17 AM

| Indicator | County Distribution | | Value | | Target | | | | |
|---|---------------------|-----|-------|-----|--------|-------|-------|-------|-----------|
| | State | US | State | US | HP2020 | Local | Trend | Score | Precision |
| rublic Water Supply - Percent of Population Served Unaffected by SDW/ | 3 | 1.5 | 1 | 1.5 | 1.5 | | 1 | 1.56 | Medium |
| ge-adjusted Heart Disease Mortality Rate per 100,000 population | 2 | 1.5 | 2 | 1.5 | 3 | | 0 | 1.50 | Medium |
| trial Fibrillation: Medicare Population | 1 | 1 | 1 | 1 | 1.5 | | 3 | 1.50 | High |
| lyperlipidemia: Medicare Population | 2 | 1 | 1 | 0 | 1.5 | | 3 | 1.50 | High |
| eople 65+ with Low Access to a Grocery Store | 1 | 2 | 1.5 | 1.5 | 1.5 | | 1.5 | 1.50 | Low |
| ercent of all Births Occurring to Teens (15-19) | 2 | 1.5 | 3 | 1.5 | 1.5 | | 0 | 1.50 | Medium |
| ercent of Births with Inadequate Birth Spacing | 2 | 1.5 | 3 | 1.5 | 1.5 | | 0 | 1.50 | Medium |
| ercent of Adults Doing Enough Physical Activity To Meet Both The Aerc | 1.5 | 1.5 | 1 | 1.5 | 2 | | 1.5 | 1.47 | Low |
| ercent of Adults Who Reported That They Always Wear a Seatbelt Whe | 1.5 | 1.5 | 1 | 1.5 | 2 | | 1.5 | 1.47 | Low |
| atio of Population to Primary Care Physicians | 1 | 1.5 | 1 | 1.5 | 1.5 | | 2 | 1.44 | Medium |
| ercent of Adults who are Overweight | 1.5 | 1.5 | 1 | 1.5 | 1.5 | | 1.5 | 1.42 | Low |
| ercent of Adults Who Reported Consuming Vegetables Less than 1 Tim | 1.5 | 1.5 | 1 | 1.5 | 1.5 | | 1.5 | 1.42 | Low |
| ercent of Adults with Doctor Diagnosed Arthritis | 1.5 | 1.5 | 1 | 1.5 | 1.5 | | 1.5 | 1.42 | Low |
| Median Household Income | 1 | 1 | 2 | 2 | 1.5 | | 1 | 1.39 | High |
| er Capita Income | 1 | 1 | 2 | 2 | 1.5 | | 1 | 1.39 | High |
| ercent of Births Where Prenatal Care began in First Trimester | 2 | 1.5 | 2 | 1.5 | 2 | | 0 | 1.39 | Medium |
| ge-adjusted Mortality Rate per 100,000 population | 2 | 1.5 | 2 | 1.5 | 1.5 | | 0 | 1.33 | Medium |
| ge-Adjusted Years of Potential Life Lost - Cerebrovascular Disease | 2 | 1.5 | 2 | 1.5 | 1.5 | | 0 | 1.33 | Medium |
| louseholds with No Car and Low Access to a Grocery Store | 1 | 1 | 1.5 | 1.5 | 1.5 | | 1.5 | 1.33 | Low |
| njury Hospital Admission Rate | 2 | 1.5 | 2 | 1.5 | 1.5 | | 0 | 1.33 | Medium |
| ife Expectancy for Males | 2 | 1 | 2 | 2 | 1.5 | | 0 | 1.33 | High |
| ercent of births Where Mother Smoked During Pregnancy | 1 | 1.5 | 2 | 1.5 | 3 | | 0 | 1.33 | Medium |
| ercentage of Screened 3-12 Grade Students w No Dental Sealants | 1 | 1.5 | 1 | 1.5 | 1.5 | | 1.5 | 1.33 | Medium |
| atio of Population to Dentist | 1 | 1.5 | 1 | 1.5 | 1.5 | | 1.5 | 1.33 | Low |
| enters Spending 30% or More of Household Income on Rent | 3 | 1 | 2 | 1 | 1.5 | | 0 | 1.33 | High |
| sthma: Medicare Population | 2 | 1 | 1 | 0 | 1.5 | | 2 | 1.28 | High |
| ongestive Heart Failure Hospital Admission Rate | 1 | 1.5 | 0 | 1.5 | 1.5 | | 2 | 1.28 | Medium |
| ercent of Adults Who are Binge Drinkers | 1.5 | 1.5 | 1 | 1.5 | 0 | | 1.5 | 1.25 | Low |
| ge-adjusted Diabetes Mortality Rate per 100,000 population | 1 | 1.5 | 1 | 1.5 | 1.5 | | 1 | 1.22 | Medium |
| ancer: Medicare Population | 1 | 2 | 1 | 1 | 1.5 | | 1 | 1.22 | High |
| ecreation and Fitness Facilities | 1 | 1 | 1.5 | 0 | 1.5 | | 2 | 1.19 | Medium |
| ge-adjusted Cerebrovascular Disease Mortality Rate per 100,000 popu | 1 | 1.5 | 1 | 1.5 | 3 | | 0 | 1.17 | Medium |
| leart Disease Hospital Admission Rate | 2 | 1.5 | 1 | 1.5 | 1.5 | | 0 | 1.17 | Medium |
| ypertension: Medicare Population | 1 | 1 | 1 | 1 | 1.5 | | 1.5 | 1.17 | High |
| ow-Income Persons who are SNAP Participants | 0 | 1 | 1.5 | 1.5 | 1.5 | | 1.5 | 1.17 | Low |
| eople 25+ with a High School Degree or Higher | 2 | 1 | 2 | 1 | 1.5 | | 0 | 1.17 | High |
| ercentage of Screened K-12 Grade Students w Obvious Dental Decay | 1 | 1.5 | 2 | 1.5 | 1.5 | | 0 | 1.17 | Medium |
| lean Travel Time to Work | 2 | 0 | 1 | 0 | 1.5 | | 2 | 1.11 | High |
| oter Turnout | 1 | 1.5 | 1 | 0 | 1.5 | | 1.5 | 1.08 | Medium |
| OPD: Medicare Population | 1 | 1 | 1 | 1 | 1.5 | | 1 | 1.06 | High |
| troke: Medicare Population | 2 | 1 | 1 | 0 | 1.5 | | 1 | 1.06 | High |
| acterial Pneumonia Hospital Admission Rate | 0 | 1.5 | 0 | 1.5 | 1.5 | | 1.5 | 1.00 | Medium |
| schemic Heart Disease: Medicare Population | 1 | 1 | 2 | 1 | 1.5 | | 0 | 1.00 | High |
| eople 65+ Living Below Poverty Level | 2 | 1 | 2 | 0 | 1.5 | | 0 | 1.00 | High |
| ge-adjusted Traffic Injury Mortality Rate per 100,000 population | 1.5 | 1.5 | 0 | 1.5 | 0 | | 1 | 0.97 | Medium |
| ccess to Exercise Opportunities | 0 | 0 | 1 | 1.5 | 1.5 | | 1.5 | 0.92 | Medium |
| olo Drivers with a Long Commute | 1 | 0 | 0 | 1.5 | 1.5 | | 1.5 | 0.92 | Medium |
| ge-Adjusted Years of Potential Life Lost - Traffic Injury | 1 | 1.5 | 0 | 1.5 | 1.5 | | 0 | 0.83 | Medium |
| ge-Adjusted Years of Potential Life Lost - Unintentional Injuries | 0 | 1.5 | 1 | 1.5 | 1.5 | | 0 | 0.83 | Medium |
| eart Failure: Medicare Population | 1 | 1 | 1 | 1 | 1.5 | | 0 | 0.83 | High |
| eople 25+ with a Bachelor's Degree or Higher | 0 | 0 | 2 | 2 | 1.5 | | 0 | 0.83 | High |
| heumatoid Arthritis or Osteoarthritis: Medicare Population | 0 | 0 | 1 | 0 | 1.5 | | 2 | 0.78 | High |
| ge-adjusted Alzheimer's Disease Mortality Rate per 100,000 population | 0 | 1.5 | 0 | 1.5 | 1.5 | | 0 | 0.67 | Medium |
| hronic Obstructive Pulmonary Disease (COPD) Hospital Admission Rate | 0 | 1.5 | 0 | 1.5 | 1.5 | | 0 | 0.67 | Medium |
| Osteoporosis: Medicare Population | 1 | 2 | 0 | 0 | 1.5 | | 0 | 0.67 | High |



County: Sample Kansas State

Total indicators: 124

Tuesday 14th of July 2015 09:31:43 AM

| Topic | Indicators | Score |
|-------------------------------------|------------|-------|
| Mental Health & Mental Disorders | 5 | 2.04 |
| Social Environment | 5 | 1.87 |
| Economy | 22 | 1.82 |
| Diabetes | 4 | 1.74 |
| Transportation | 7 | 1.73 |
| Maternal, Fetal & Infant Health | 10 | 1.70 |
| Environment | 15 | 1.67 |
| Exercise, Nutrition, & Weight | 20 | 1.67 |
| Prevention & Safety | 8 | 1.67 |
| Immunizations & Infectious Diseases | 4 | 1.63 |
| Substance Abuse | 6 | 1.61 |
| Mortality Data | 26 | 1.58 |
| Children's Health | 5 | 1.57 |
| Wellness & Lifestyle | 4 | 1.57 |
| Cancer | 3 | 1.52 |
| Public Safety | 7 | 1.50 |
| Older Adults & Aging | 21 | 1.42 |
| Education | 4 | 1.41 |
| Other Chronic Diseases | 5 | 1.39 |
| Heart Disease & Stroke | 14 | 1.35 |
| Respiratory Diseases | 6 | 1.31 |
| Oral Health | 3 | 1.28 |

Scoring Results by Topic

County: Sample Kansas State

Topic Selection:

Total indicators: 124

Tuesday 14th of July 2015 09:30:17 AM

Choose a topic here ↓

2.04

Mental Health & Mental Disorders

Topic Score:

After pasting in scoring results, refresh the Excel workbook to see comparison and indicator scores:

In Excel: Data menu --> Connections --> Refresh All

NOTE: If using Excel for Mac, this action must be completed in the Excel window (Data --> Refresh) and not in the menu bar at the top of the your screen.

Mental Health & Mental Disorders in Sample County

| | County Distribution | | Value | | Target | | | | |
|--|---------------------|-----|-------|-----|--------|-------|-------|-------|-----------|
| Indicator | State | US | State | US | HP2020 | Local | Trend | Score | Precision |
| Depression: Medicare Population | 3 | 3 | 2 | 3 | 1.5 | | 3 | 2.67 | High |
| Age-Adjusted Years of Potential Life Lost - Alzheimers | 3 | 1.5 | 2 | 1.5 | 1.5 | | 3 | 2.17 | Medium |
| Alzheimer's Disease or Dementia: Medicare Population | 2 | 3 | 2 | 2 | 1.5 | | 1 | 1.89 | High |
| Age-Adjusted Years of Potential Life Lost - Suicide | 2 | 1.5 | 2 | 1.5 | 1.5 | | 2 | 1.78 | Medium |
| Age-adjusted Suicide Mortality Rate per 100,000 population | 1.5 | 1.5 | 1 | 1.5 | 3 | | 2 | 1.69 | Medium |